

SNUTP 01-040

DESY 01-194

November, 2001

Polarization Effects in $B \rightarrow K_1(1270) + J/\psi$ Decays^(a)G. Kramer, ^(b)H. S. Song, and ^(b)Chaehyun Yu.^(a) *II. Institut für Theoretische Physik, Universität Hamburg, D-22761 Hamburg, Germany*^(b) *Center for Theoretical Physics and School of Physics, Seoul National University,
Seoul 151-742, Korea***Abstract**

The joint angular distribution of the decay $B \rightarrow J/\psi V$, where V is an axial vector or vector resonance, followed by the subsequent decay processes of the J/ψ and V is calculated using the covariant density method. In particular, the case where V is the axial vector meson $K_1(1270)$ which decays into $K\rho$ is considered as well as the case that V is the vector meson $K^*(890)$ which decays into $K\pi$.

Typeset using REVTeX

I. INTRODUCTION

Recently the decays $B^0 \rightarrow K_1^0(1270)J/\psi$ and $B^+ \rightarrow K_1^+(1270)J/\psi$ have been observed in the Belle detector at the KEKB asymmetric e^+e^- collider with significant branching ratios of the order of 10^{-3} [1]. So these decays might be useful for CP violation studies similar to the decay $B \rightarrow K^*(892)J/\psi$. The final state in the decay $B \rightarrow K_1J/\psi$ is a mixture of $CP = \pm 1$ eigenstates depending on the angular momentum of the K_1 and J/ψ . For CP violation studies, the relative strengths of the two eigenstates must be known. These can be determined from an analysis of the joint angular correlation of the decay products of $K_1 \rightarrow K\rho$ and $J/\psi \rightarrow l^+l^-$, where l may be an electron or muon, respectively [2]. Since the branching fraction of the decay $K_1 \rightarrow K\rho$ is quite large about 42% [3] and the decay $J/\psi \rightarrow l^+l^-$ has a rather distinct signature, which easily can be measured, it is to be expected that the full angular correlation will be measured as soon as a large data sample has been collected at Belle and at BABAR. In fact, this has been achieved recently for the decay $B \rightarrow K^*J/\psi$ with $K^* \rightarrow K\pi$ and $J/\psi \rightarrow l^+l^-$ with high precision at BABAR [4] and Belle [5] improving earlier measurements by the CLEO [6] and CDF [7] collaborations.

The joint angular distribution of a B meson decaying into the vector (or axial vector) particles V_1 and V_2 with subsequent decays of V_1 and V_2 , respectively, is described by three angles. In the original derivation of the angular correlation for the decay $B \rightarrow K^*J/\psi$ with $K^* \rightarrow K\pi$ and $J/\psi \rightarrow l^+l^-$ [8] the usual polar and azimuthal angles for the decays of K^* and J/ψ were used, describing the decay matrix elements of the decay products for B , K^* and J/ψ in the familiar helicity basis [9]. The CP parity information, however, is more easily extracted by introducing the B decay amplitudes A_0 , $A_{\parallel}$ and $A_{\perp}$ in the transversity basis [2]. In this basis, using in the transversity system angles θ_{tr} and ϕ_{tr} , defined as polar and azimuthal angles of the l^+ in the J/ψ rest frame, the angular correlation was presented in [10]. Following the earlier work the angular correlation for the decay chain $B \rightarrow K_1J/\psi$, $K_1 \rightarrow K\rho$ and $J/\psi \rightarrow l^+l^-$ can be derived as well either in the helicity or the transversity

basis. Since the matrix elements of the two decay processes $B \rightarrow K^* J/\psi$ and $B \rightarrow K_1 J/\psi$ have the same Lorentz covariant form, although the invariant form factors can be different in size and CP properties, it is more convenient to apply the method of covariant density matrices [11] instead. Then it is quite simple to incorporate the fact that the two subsequent decays $K^* \rightarrow K\pi$ and $K_1 \rightarrow K\rho$ produce different contributions in the decay density matrix of the J/ψ .

In this short note we derive the angular distribution of the decay $J/\psi \rightarrow l^+ l^-$ in terms of the density matrix $\rho_{\lambda\lambda'}$. The explicit form of $\rho_{\lambda\lambda'}$ for the two cases, $B \rightarrow K_1(1270) J/\psi$ and $B \rightarrow K^*(892) J/\psi$, is calculated covariantly. With this result the full angular distribution for both decays are derived in the transversity basis. It is clear that this method is quite general and can be applied to any B decay into two spin-1 particles (either vector or axial vector) and their subsequent decays.

II. DECAY OF $J/\psi \rightarrow L^+ L^-$

Neglecting weak effects the process $J/\psi(q) \rightarrow l^+(q_1) l^-(q_2)$ can be described by the amplitude

$$\mathcal{M} = \frac{f_1}{2} \epsilon^\mu(q) \bar{u}(q_2) \gamma_\mu v(q_1), \quad (1)$$

where f_1 is a coupling constant. Then the decay distribution of the J/ψ with polarization vector $\epsilon^\mu(q)$ after it is produced in any process is obtained from

$$|\mathcal{M}|^2 = |f_1|^2 \langle \epsilon_\mu(q) \epsilon_\nu^*(q) \rangle (q_1^\mu q_2^\nu + q_1^\nu q_2^\mu - \frac{q^2}{2} g^{\mu\nu}), \quad (2)$$

where the lepton polarizations have been summed. Here, the ensemble averaged value $\langle \epsilon_\mu(q) \epsilon_\nu^*(q) \rangle$ can be replaced by the covariant density matrix $\rho_{\mu\nu}$ of the J/ψ which is obtained explicitly from the B decay process into the J/ψ . The most general form of $\rho^{\mu\nu}$ can be written as [11]

$$\rho_{\mu\nu} = \frac{1}{3}(-g^{\mu\nu} + \frac{q_\mu q_\nu}{m_1^2}) - \frac{i}{2m_1}\epsilon_{\mu\nu\sigma\tau}q^\sigma\mathcal{P}^\tau - \frac{1}{2}\mathcal{Q}_{\mu\nu}, \quad (3)$$

where the polarization vector $\mathcal{P}^\tau$ and the polarization tensor $\mathcal{Q}^{\mu\nu}$ can be obtained explicitly. The covariant density matrix $\rho^{\mu\nu}$ is related to the density matrix $\rho_{\lambda\lambda'}$ in the spin momentum basis as follows

$$\rho^{\mu\nu} = \epsilon^\mu(q, \lambda)\rho_{\lambda\lambda'}\epsilon^{\nu*}(q, \lambda'). \quad (4)$$

Here the projection operation $\epsilon^\mu(q, \lambda)\epsilon^{\nu*}(q, \lambda')$ is given by [11]

$$\epsilon^\mu(q, \lambda)\epsilon^{\nu*}(q, \lambda') = \frac{1}{3}\left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{m_1^2}\right)\delta_{\lambda'\lambda} - \frac{i}{2m_1}\epsilon^{\mu\nu\sigma\tau}q_\sigma n_\tau^i (S^i)_{\lambda'\lambda} - \frac{1}{2}n_i^\mu n_j^\nu (S^{ij})_{\lambda'\lambda}, \quad (5)$$

where the (S^i) are the standard matrix representations of spin-1 angular momentum operators and (S^{ij}) are traceless and symmetric matrices defined as

$$S^{ij} = S^i S^j + S^j S^i - \frac{4}{4}\delta^{ij}I. \quad (6)$$

In Eq. (5), $(n_0^\mu, n_1^\mu, n_2^\mu, n_3^\mu)$ form a tetrad with $n_0^\mu = (E/m, \vec{q}/m)$ and n_3^μ is the Lorentz boost from $\hat{q}$ in the J/ψ rest frame [11]. Then $\mathcal{P}^\mu$ and $\mathcal{Q}^{\mu\nu}$ in Eq. (3) can be written as

$$\begin{aligned} \mathcal{P}^\mu &= n_i^\mu \text{Tr}(S^i \rho) \\ \mathcal{Q}^{\mu\nu} &= n_i^\mu n_j^\nu \text{Tr}(S^{ij} \rho). \end{aligned} \quad (7)$$

Then Eq. (2) becomes

$$|\mathcal{M}|^2 \sim |f|^2 m_1^2 \rho_{\lambda\lambda'} \left[\frac{1}{3}\delta_{\lambda'\lambda} + \frac{1}{m_1^2} q_1 \cdot n^i q_1 \cdot n^j (S^{ij})_{\lambda'\lambda} \right], \quad (8)$$

In the transversity basis, defined as in Fig. 1 in the J/ψ rest frame, it is

$$\begin{aligned} d\Gamma \sim & \left[1 + \rho_{00} + (1 - 3\rho_{00}) \sin^2 \theta_{tr} \cos^2 \phi_{tr} \right. \\ & + 2\text{Re}(\rho_{1-1})(\sin^2 \theta_{tr} \sin^2 \phi_{tr} - \cos^2 \theta_{tr}) \\ & - 2\text{Im}(\rho_{1-1}) \sin^2 \theta_{tr} \sin 2\phi_{tr} \\ & + \sqrt{2}\text{Re}(\rho_{10} - \rho_{-11}) \sin 2\theta_{tr} \cos \phi_{tr} \\ & \left. - \sqrt{2}\text{Im}(\rho_{10} + \rho_{-11}) \sin 2\theta_{tr} \sin \phi_{tr} \right], \end{aligned} \quad (9)$$

where (θ_{tr}, ϕ_{tr}) are the polar and azimuthal angles of the outgoing lepton l^+ in the transversity basis as defined in [2,10] where $(\hat{n}_3, \hat{n}_1, \hat{n}_2)$ is along the $(\hat{x}, \hat{y}, \hat{z})$ axis.

III. DENSITY MATRIX OF J/ψ

The explicit values of the density matrix elements $\rho_{\lambda\lambda'}$ are calculated from the amplitude of the J/ψ production process. The matrix element of the $B \rightarrow J/\psi V$ (V is either K_1 or K^*) decay is given in terms of three independent Lorentz scalars A , B and C as [12,8]

$$\mathcal{M} = A\epsilon_1^* \cdot \epsilon_2^* + \frac{B}{m_1 m_2} \epsilon_1^* \cdot q \epsilon_2^* \cdot k + \frac{iC}{m_1 m_2} \epsilon^{\mu\nu\sigma\tau} \epsilon_{1\mu}^* \epsilon_{2\nu}^* q_\sigma k_\tau, \quad (10)$$

where $\epsilon_i (i = 1, 2)$ are the polarization vectors of J/ψ and V , respectively, and m_2 and k^μ are the mass and momentum of the V particle. Instead of the scalar amplitudes a , b and c it is more convenient to introduce the three transversity amplitudes A_0 , $A_\parallel$ and $A_\perp$ if one wants to specify the CP property of the decay process. They are related to A, B, C by [10]

$$\begin{aligned} A_0 &= -xA - (x^2 - 1)B, \\ A_\parallel &= \sqrt{2} A, \\ A_\perp &= \sqrt{2(x^2 - 1)} C, \end{aligned} \quad (11)$$

where x is defined as

$$x = \frac{k \cdot q}{m_1 m_2} = \frac{m_B^2 - m_1^2 - m_2^2}{2m_1 m_2}. \quad (12)$$

When the decay of the B into J/ψ and V is specified by the polarization vectors $\epsilon_1(q, \lambda_1)$ and $\epsilon_2(k, \lambda_2)$, respectively, the density matrix elements of the J/ψ and V final states can be obtained from the square of Eq. (10) and by using Eq. (5) for each $\epsilon_i (i = 1, 2)$ as follows,

$$\begin{aligned} \rho_{\lambda_1 \lambda'_1, \lambda_2 \lambda'_2} &\sim \frac{1}{9} (|A_0|^2 + |A_\parallel|^2 + |A_\perp|^2) \delta_{\lambda_1 \lambda'_1, \lambda_2 \lambda'_2} \\ &- \frac{2}{3\sqrt{\Delta}} \text{Re}(A_\parallel^* A_\perp) \{ m_1 n_1^i \cdot k (S^i)_{\lambda_1 \lambda'_1} \delta_{\lambda_2 \lambda'_2} + m_2 n_2^k \cdot q \delta_{\lambda_1 \lambda'_1} (S^k)_{\lambda_2 \lambda'_2} \} \\ &+ \frac{1}{\Delta} \{ (|A_\parallel|^2 + |A_\perp|^2) m_1 m_2 n_1^i \cdot k n_2^k \cdot q \\ &\quad + \sqrt{\frac{\Delta}{2}} \text{Im}(A_0^* A_\perp) \langle q k n_1^i n_2^k \rangle \\ &\quad - \frac{\sqrt{2}}{4} \Delta \text{Re}(A_0^* A_\parallel) (n_1^i \cdot n_2^k - \frac{4q \cdot k}{\Delta} n_1^i \cdot k n_2^k \cdot q) \} (S^i)_{\lambda_1 \lambda'_1} (S^k)_{\lambda_2 \lambda'_2} \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{3\Delta}(2|A_0|^2 - |A_{\parallel}|^2 - |A_{\perp}|^2)\left\{ m_1^2 n_1^i \cdot k n_1^j \cdot k (S^{ij})_{\lambda_1 \lambda'_1} \delta_{\lambda_2 \lambda'_2} \right. \\
& \quad \left. + m_2^2 n_2^k \cdot q n_2^l \cdot q \delta_{\lambda_1 \lambda'_1} (S^{kl})_{\lambda_2 \lambda'_2} \right\} \\
& -\frac{1}{\Delta}\left\{ \sqrt{2}\text{Im}(A_0^* A_{\parallel}) \langle qk n_1^i n_2^k \rangle \right. \\
& \quad + \frac{2}{\sqrt{\Delta}}\text{Re}(A_{\parallel}^* A_{\perp}) m_1 m_2 n_1^i \cdot k n_2^k \cdot q \\
& \quad - \sqrt{\frac{\Delta}{2}}\text{Re}(A_0^* A_{\perp}) \left(n_1^i \cdot n_2^k - \frac{4q \cdot k}{\Delta} n_1^i \cdot k n_2^k \cdot q \right) \left. \right\} \\
& \quad \times \{ m_2 n_2^l \cdot q (S^i)_{\lambda_1 \lambda'_1} (S^{kl})_{\lambda_2 \lambda'_2} + m_1 n_1^j \cdot q (S^{ij})_{\lambda_1 \lambda'_1} (S^k)_{\lambda_2 \lambda'_2} \} \\
& + \left\{ \frac{4}{\Delta^2} |A_0|^2 m_1^2 m_2^2 n_1^i \cdot k n_1^j \cdot k n_2^k \cdot q n_2^l \cdot q \right. \\
& \quad + \frac{1}{8} |A_{\parallel}|^2 \left(n_1^i \cdot n_2^k - \frac{4q \cdot k}{\Delta} n_1^i \cdot k n_2^k \cdot q \right) \left(n_1^j \cdot n_2^l - \frac{4q \cdot k}{\Delta} n_1^j \cdot k n_2^l \cdot q \right) \\
& \quad + \frac{1}{2\Delta} |A_{\perp}|^2 \langle qk n_1^i n_2^k \rangle \langle qk n_1^j n_2^l \rangle \\
& \quad - \frac{\sqrt{2}}{2\Delta} \text{Re}(A_0^* A_{\parallel}) m_1 m_2 n_1^i \cdot k n_2^k \cdot q \left(n_1^j \cdot n_2^l - \frac{4q \cdot k}{\Delta} n_1^j \cdot k n_2^l \cdot q \right) \\
& \quad + \frac{2\sqrt{2}}{\Delta\sqrt{\Delta}} \text{Im}(A_0^* A_{\perp}) m_1 m_2 n_1^i \cdot k n_2^k \cdot q \langle qk n_1^j n_2^l \rangle \\
& \quad \left. - \frac{1}{2\sqrt{\Delta}} \text{Im}(A_{\parallel}^* A_{\perp}) \left(n_1^i \cdot n_2^k - \frac{4q \cdot k}{\Delta} n_1^i \cdot k n_2^k \cdot q \right) \langle qk n_1^j n_2^l \rangle \right\} (S^{ij})_{\lambda_1 \lambda'_1} (S^{kl})_{\lambda_2 \lambda'_2}. \quad (13)
\end{aligned}$$

Here the density matrix is multiplied by $(|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2)$ for convenience and Δ and $\langle qk n_1^j n_2^l \rangle$ are defined as

$$\begin{aligned}
\Delta &= (m_B^4 + m_1^4 + m_2^4 - 2m_B^2 m_1^2 - 2m_B^2 m_2^2 - 2m_1^2 m_2^2) \\
&= 4[(q \cdot k)^2 - m_1^2 m_2^2] = 4m_1^2 m_2^2 (x^2 - 1), \quad (14)
\end{aligned}$$

$$\langle qk n_1^j n_2^l \rangle = \epsilon^{\mu\nu\sigma\tau} q_{\mu} k_{\nu} (n_{\sigma}^j)_1 (n_{\tau}^l)_2. \quad (15)$$

Now the density matrix element of J/ψ , after V decays, can be obtained by specifying the amplitude of the V particle decay process. If the explicit form of the V decay process is given, one can explicitly obtain the density matrix of the V , $\rho_{\lambda'_2 \lambda_2}^D(V)$, for its decay process. Then the density matrix of the J/ψ can be obtained from the product

$$\rho_{\lambda_1 \lambda'_1, \lambda_2 \lambda'_2} \rho_{\lambda'_2 \lambda_2}^D(V). \quad (16)$$

The amplitude of the decay process $K_1(k, \lambda_2) \rightarrow K(\tilde{k})\rho(\tilde{k}')$ is described by

$$\mathcal{M}_2 = a\epsilon_2 \cdot \epsilon_3^* + \frac{b}{m_2 m_\rho} \epsilon_2 \cdot \tilde{k}' \epsilon_3^* \cdot k, \quad (17)$$

where ϵ_3^μ and $\tilde{k}'$ are the polarization vector and momentum of ρ . Equation (13) can be used for the decay process $K_1 \rightarrow K\rho$ if some notations are changed, and after the polarization of ρ is summed over, we obtain

$$\rho_{\lambda'_2 \lambda_2}^D(K_1) = \frac{1}{3} \delta_{\lambda'_2 \lambda_2} - \frac{2m_2^2}{\tilde{\Delta}} (1 - \xi) \tilde{k} \cdot n_2^k \tilde{k} \cdot n_2^l (S_2^{kl})_{\lambda'_2 \lambda_2}, \quad (18)$$

where $\tilde{\Delta}$ and ξ are defined as

$$\tilde{\Delta} = (m_2^4 + m_K^4 + m_\rho^4 - 2m_2^2 m_K^2 - 2m_2^2 m_\rho^2 - 2m_K^2 m_\rho^2), \quad (19)$$

$$\xi = \frac{3|a_\parallel|^2}{2(|a_0|^2 + |a_\parallel|^2)}. \quad (20)$$

The linear polarization amplitudes in the K_1 decay are related to the form factors in Eq. (17) and helicity amplitudes H_0 and $H_\pm$ as

$$a_0 = -\tilde{x}a - (\tilde{x}^2 - 1)b = H_0, \quad (21)$$

$$a_\parallel = \sqrt{2}a = (H_{+1} + H_{-1})/\sqrt{2} = \sqrt{2}H_+, \quad (22)$$

where $\tilde{x} = (m_2^2 - m_K^2 + m_\rho^2)/2m_2 m_\rho$.

Likewise, the amplitude of the decay process $K^*(k, \lambda_2) \rightarrow K(\tilde{k})\pi(\tilde{k}')$ is given by

$$\mathcal{M}_3 = f_3 \epsilon^\mu(k, \lambda_2) (\tilde{k} - \tilde{k}')_\mu = 2f_3 \epsilon^\mu(k, \lambda_2) \tilde{k}_\mu, \quad (23)$$

and the density matrix of the decay process becomes

$$\rho_{\lambda'_2 \lambda_2}^D(K^*) = \frac{1}{3} \delta_{\lambda'_2 \lambda_2} - \frac{2m_2^2}{\tilde{\Delta}} \tilde{k} \cdot n_2^k \tilde{k} \cdot n_2^l (S_2^{kl})_{\lambda'_2 \lambda_2}, \quad (24)$$

where $\tilde{\Delta}$ is defined in the same way as in Eq. (19) after replacing the subscripts K, ρ by K, π . We see that the density matrix for K^* to $K\pi$ is obtained from that of K_1 to $K\rho$ in Eq. (18) if we replace $\xi = 0$.

Considering the two processes simultaneously, we write the decay density matrix elements of the two V particles (K_1 and K^*), Eqs. (18) and (24), as

$$\rho_{\lambda'_2\lambda_2}^D = \frac{1}{3}\delta_{\lambda'_2\lambda_2} + D\tilde{k}\cdot n_2^k \tilde{k}\cdot n_2^l (S^{kl})_{\lambda'_2\lambda_2}, \quad (25)$$

and then we have

$$\begin{aligned} D &= -\frac{2m_2^2}{\tilde{\Delta}}, & \text{for } K^* \rightarrow K\pi, \\ &= -\frac{2m_2^2}{\tilde{\Delta}}(1 - \xi), & \text{for } K_1 \rightarrow K\rho. \end{aligned} \quad (26)$$

The explicit form of the density matrix elements of J/ψ neglecting the subscript 1 in λ_1 and n_1 is now obtained as follows,

$$\begin{aligned} \rho_{\lambda\lambda'} &\sim \rho_{\lambda_1\lambda'_1,\lambda_2\lambda'_2}\rho_{\lambda'_2\lambda_2}^D \\ &= \frac{1}{3}\delta_{\lambda\lambda'} \left[\frac{1}{3}(|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2) \right. \\ &\quad \left. + \frac{D\tilde{\Delta}}{12m_2^2} \left\{ 1 - \frac{48m_2^4}{\Delta\tilde{\Delta}}(q\cdot J)^2 \right\} (2|A_0|^2 - |A_{\parallel}|^2 - |A_{\perp}|^2) \right] \\ &\quad + (S^i)_{\lambda\lambda'} \left[-\frac{2}{3\sqrt{\Delta}} \text{Re}(A_{\parallel}^* A_{\perp}) m_1 k\cdot n^i \right. \\ &\quad \left. + \frac{2\sqrt{2}m_2}{\Delta} D \left\{ 2\text{Im}(A_0^* A_{\parallel}) q\cdot J \langle qk\tilde{k}n^i \rangle \right. \right. \\ &\quad \left. + \frac{m_1}{24} \sqrt{2\Delta} \Delta \text{Re}(A_{\parallel}^* A_{\perp}) \left\{ 1 - \frac{48m_2^4}{\Delta\tilde{\Delta}}(q\cdot J)^2 \right\} k\cdot n^i \right. \\ &\quad \left. \left. + \sqrt{\Delta} \text{Re}(A_0^* A_{\perp}) q\cdot J (J\cdot n^i - \frac{4q\cdot k q\cdot J}{\Delta} k\cdot n^i) \right\} \right] \\ &\quad + (S^{ij})_{\lambda\lambda'} \left[-\frac{m_1^2}{3\Delta} \left(1 + \frac{D\tilde{\Delta}}{2m_2^2} \right) (2|A_0|^2 - |A_{\parallel}|^2 - |A_{\perp}|^2) k\cdot n^i k\cdot n^j \right. \\ &\quad \left. + D \left\{ \frac{16m_1^2 m_2^2}{\Delta^2} (q\cdot J)^2 k\cdot n^i k\cdot n^j |A_0|^2 \right. \right. \\ &\quad \left. + \frac{1}{2} |A_{\parallel}|^2 (J\cdot n^i - \frac{4q\cdot k q\cdot J}{\Delta} k\cdot n^i) (J\cdot n^j - \frac{4q\cdot k q\cdot J}{\Delta} k\cdot n^j) \right. \\ &\quad \left. + \frac{2}{\Delta} |A_{\perp}|^2 \langle qk\tilde{k}n^i \rangle \langle qk\tilde{k}n^j \rangle \right. \\ &\quad \left. - \frac{4\sqrt{2}}{\Delta} m_1 m_2 (q\cdot J) \text{Re}(A_0^* A_{\parallel}) k\cdot n^i (J\cdot n^j - \frac{4q\cdot k q\cdot J}{\Delta} k\cdot n^j) \right. \\ &\quad \left. - \frac{8\sqrt{2}}{\Delta\sqrt{\Delta}} m_1 m_2 (q\cdot J) \text{Im}(A_0^* A_{\perp}) k\cdot n^i \langle qk\tilde{k}n^j \rangle \right. \\ &\quad \left. \left. + \frac{2}{\sqrt{\Delta}} \text{Im}(A_{\parallel}^* A_{\perp}) (J\cdot n^i - \frac{4q\cdot k q\cdot J}{\Delta} k\cdot n^i) \langle qk\tilde{k}n^j \rangle \right\} \right], \quad (27) \end{aligned}$$

where $J = -\tilde{k} + \frac{k\cdot\tilde{k}}{m_2^2}k$. This density matrix is manifestly covariant and from the explicit form of matrices $(S^i), (S^{ij})$, one can obtain the angular distribution of $B \rightarrow K_1 J/\psi$ as well as that

of $B \rightarrow K^* J/\psi$. The angular distribution for the decay process $B \rightarrow K_1(1270) J/\psi$ ($K_1 \rightarrow K\rho, J/\psi \rightarrow l^+ l^-$) in the transversity basis is derived from Eqs. (9) and (27):

$$\begin{aligned} \frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_1 d\cos\theta_{tr} d\phi_{tr}} = \frac{9}{32\pi} \frac{1}{1+3\alpha} & \left[2|A_0(t)|^2 \left\{ (\cos^2\theta_1 + \alpha)(1 - \sin^2\theta_{tr} \cos^2\phi_{tr}) \right\} \right. \\ & + |A_{\parallel}(t)|^2 \left\{ \sin^2\theta_1(1 - \sin^2\theta_{tr} \sin^2\phi_{tr}) \right. \\ & \quad \left. + \alpha(1 + \sin^2\theta_{tr} \cos^2\phi_{tr}) \right\} \\ & + |A_{\perp}(t)|^2 \left\{ \sin^2\theta_1 \sin^2\theta_{tr} + \alpha(1 + \sin^2\theta_{tr} \cos^2\phi_{tr}) \right\} \\ & + \text{Im}(A_{\parallel}^*(t) A_{\perp}(t)) \sin^2\theta_1 \sin 2\theta_{tr} \sin\phi_{tr} \\ & - \frac{1}{\sqrt{2}} \text{Re}(A_0^*(t) A_{\parallel}(t)) \sin 2\theta_1 \sin^2\theta_{tr} \sin 2\phi_{tr} \\ & \left. + \frac{1}{\sqrt{2}} \text{Im}(A_0^*(t) A_{\perp}(t)) \sin 2\theta_1 \sin 2\theta_{tr} \cos\phi_{tr} \right], \quad (28) \end{aligned}$$

where θ_1 is the angle between the K_1 direction and the K direction in the K_1 rest frame and the three amplitudes are normalized as $|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2 = 1$. Here, α is defined as

$$\alpha = \frac{1}{3} \frac{\xi}{1-\xi} = \frac{|a_{\parallel}|^2}{2|a_0|^2 - |a_{\parallel}|^2}. \quad (29)$$

The angular distribution in the process $B \rightarrow K^* J/\psi$ ($K^* \rightarrow K\pi, J/\psi \rightarrow l^+ l^-$) in the transversity basis can be obtained by putting $\xi = 0$ or $\alpha = 0$ in Eq. (28). The angular correlation for $B \rightarrow K_1 J/\psi$ depends also on details of the decay $K_1 \rightarrow K\rho$ in contrast to $K^* \rightarrow K\pi$ and this is related to the fact that the decay $K_1 \rightarrow K\rho$ depends on two independent matrix elements. In order to analyze the transversity amplitudes for the B decay, one must know the ratio of three decay matrix elements for $K_1 \rightarrow K\rho$ or otherwise one must determine them from the angular correlation together with the B decay transversity amplitudes.

ACKNOWLEDGMENTS

The authors would like to thank Prof. S. K. Kim for her suggestion to derive the new angular distribution. The work is supported by the DFG-KOSEF Collaboration (2000) under

the contracts 446 KOR-113/137/0-1 (DFG) and 20005-111-02-2 (KOSEF). H. S. Song and G. Kramer thank their guest institutes for the warm hospitality during visits in the summer and fall of 2001.

REFERENCES

- [1] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **87**, 161601 (2001).
- [2] I. Dunietz, H. J. Lipkin, H. Quinn, A. Snider and W. Toki, Phys. Rev. D **43**, 2193 (1991).
- [3] D. E. Groom *et al.* [Particle Data Group Collaboration], Eur. Phys. J. C. **15**, 1 (2000).
- [4] BABAR Collaboration, B. Aubert *et al.*, BABAR-PUB-01/05, SLAC-PUB-8898, July 20, 2001.
- [5] Belle Collaboration, K. Abe *et al.*, BELLE-CONF-0105 (2001).
- [6] CLEO Collaboration, C. P. Jessop *et al.*, Phys. Rev. Lett. **79**, 4533 (1997).
- [7] CDF Collaboration, T. Affolder *et al.*, Phys. Rev. Lett. **84** 4668 (2000).
- [8] G. Kramer and W. F. Palmer, Phys. Rev. D **45**, 193 (1992).
- [9] M. Jacob and G. C. Wick, Ann. Phys. **7**, 404 (1959).
- [10] A. S. Dighe, I. Dunietz, H. J. Lipkin and J. L. Rosner, Phys. Lett. B **369**, 144 (1996) [hep-ph/9511363].
- [11] S. Y. Choi, Taeyeon Lee and H. S. Song, Phys. Rev. D **40**, 2477 (1989).
- [12] G. Valencia, Phys. Rev. D **39**, 3339 (1989).

Figure Captions

Fig. 1 : The definitions of the angles in the transversity basis. θ_{tr} and ϕ_{tr} are polar and azimuthal angles of l^+ in the J/ψ rest frame. θ_1 is the angle between the K_1 direction and the K direction in the K_1 rest frame.

FIGURES




FIG. 1.